

ERRATUM

2018 대한부정맥학회 비판막성 심방세동 환자의 뇌졸중 예방 지침

¹경희대학교병원 순환기내과, ²연세대학교 세브란스병원 순환기내과, ³서울대학교병원 순환기내과,
⁴서울대학교 분당병원 순환기내과, ⁵서울대학교 보라매병원 순환기내과, ⁶중앙대학교병원 순환기내과,
⁷고려대학교 구로병원 순환기내과, ⁸성균관대학교 삼성서울병원 순환기내과

이정명¹ · 정보영² · 차명진³ · 이지현⁴ · 임우현⁵ · 김태훈² · 신승용⁶ · 엄재선² · 임홍익⁷ · 김진배¹ · 김준수⁸

2018 KHRS Guidelines for Stroke Prevention Therapy in Korean Patients with Nonvalvular Atrial Fibrillation

Jung Myung Lee¹, Boyoung Joung², Myung-Jin Cha³, Ji Hyun Lee⁴, Woo Hyun Lim⁵, Tae-Hoon Kim², Seung Yong Shin⁶,
 Jae-Sun Uhm², Hong Euy Lim⁷, Jin-Bae Kim¹, and Jun Soo Kim⁸

¹Division of Cardiology, Kyung Hee University Hospital, Seoul; ²Division of Cardiology, Department of Internal Medicine, Severance Cardiovascular Hospital, Yonsei University College of Medicine, Seoul; ³Department of Internal Medicine, Seoul National University Hospital, Seoul; ⁴Division of Cardiology, Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam; ⁵Division of Cardiology, Department of Internal Medicine, Boramae Medical Center, Seoul National University College of Medicine, Seoul; ⁶Department of Cardiology, Chung-Ang University Hospital, Seoul; ⁷Cardiovascular Center, Korea University Guro Hospital, Seoul; ⁸Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

Korean J Med 2018;93:87-109. <https://doi.org/10.3904/kjm.2018.93.2.87>

다음과 같이 본문 내 Table 5에 오류가 있어 수정합니다.

Table 5. Guidance for temporary interruption of NOAC before surgery

	Dabigatran	Apixaban, edoxaban, rivaroxaban	Low-risk surgery	High-risk surgery
	Low-risk surgery	High-risk surgery		
Creatinine clearance >80 mL/min	24 hours	48 hours	24 hours	48 hours
Creatinine clearance 50-80 mL/min	36 hours	72 hours	24 hours	48 hours
Creatinine clearance 30-50 mL/min	48 hours	96 hours	24 hours	48 hours
Creatinine clearance 15-30 mL/min	Not indicated	Not indicated	36 hours	48 hours

NOAC, non-vitamin K antagonist oral anticoagulant.

Table 5. Guidance for temporary interruption of NOAC before surgery

	Dabigatran		Apixaban, edoxaban, rivaroxaban	
	Low risk surgery	High risk surgery	Low risk surgery	High risk surgery
Creatinine clearance > 80 mL/min	24 hours	48 hours	24 hours	48 hours
Creatinine clearance 50- 80 mL/min	36 hours	72 hours	24 hours	48 hours
Creatinine clearance 30-50 mL/min	48 hours	96 hours	24 hours	48 hours
Creatinine clearance 15-30 mL/min	Not indicated	Not indicated	36 hours	48 hours

NOAC, non-vitamin K antagonist oral anticoagulant.