## 혈액투석 환자에서 Bioelectrical Impedance Analysis를 활용하여 측정한 과수분량과 근육량 감소와 사망률의 상관관계

<sup>1</sup>차의과학대학교 분당차병원 내과, <sup>2</sup>용인세브란스병원 내과


김은주<sup>1</sup> · 서상오<sup>1</sup> · 최유범<sup>1</sup> · 이미정<sup>1</sup> · 이정은<sup>2</sup> · 김형종<sup>1</sup>

## Relative Association of Overhydration and Muscle Wasting with Mortality in Hemodialysis Patients: Assessment by Bioelectrical Impedance Analysis


Eunju Kim<sup>1</sup>, Sang Oh Seo<sup>1</sup>, Yu Bum Choi<sup>1</sup>, Mi Jung Lee<sup>1</sup>, Jeong Eun Lee<sup>2</sup>, and Hyung Jong Kim<sup>1</sup>

<sup>1</sup>Department of Internal Medicine, CHA Bundang Medical Center, CHA University, Seongnam; <sup>2</sup>Department of Internal Medicine, Yongin Severance Hospital, Yongin, Korea

Korean J Med 2018;93:548-555. https://doi.org/10.3904/kjm.2018.93.6.548 다음과 같이 본문 내 Figure 1에 오류가 있어 수정합니다.



**Figure 1.** Kaplan-Meier survival curve of the two groups for all-cause mortality. ECW, extracellular water; TBW, total body water.



**Figure 1.** Kaplan-Meier survival curve of the two groups for all-cause mortality. ECW, extracellular water; TBW, total body water.

## Copyright © 2019 The Korean Association of Internal Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/3.0/) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.